

September 1, 2015



# Math 3012 - Applied Combinatorics Lecture 5

William T. Trotter  
trotter@math.gatech.edu

# Multinomial Coefficients

**Problem** How many different arrangements of

AABBCCCCCDEEEEEEEFFFFFFF ?

**Answer**

$$\binom{26}{2,3,1,6,6,8} = \frac{26!}{2! 3! 1! 6! 6! 8!}$$

**Note** Informally, this is known as the "MISSISSIPPI" problem.

# Binomial and Multinomial Coefficients

## Observation

When there are only two parts, a multinomial coefficient is just a binomial coefficient. So for example,

$$\binom{26}{7,19} = \binom{26}{7}$$

**However** You should only use the binomial notation in this case.

# The Binomial Theorem

## Theorem

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$$

## Explanation

$$(x + y)^n = (x + y)(x + y)(x + y)(x + y) \dots (x + y)$$

From each of  $n$  terms, you either take  $x$  or  $y$ , so if  $k$  is the number of times you take  $y$ , then you take  $x$  exactly  $n - k$  times.

# Applying the Binomial Theorem

## Theorem

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$$

**Problem** What is the coefficient of  $a^{14}b^{18}$  in

$$(3a^2 - 5b)^{25}$$

**Answer**

$$\binom{25}{7} 3^7 (-5)^{18}$$

# The Multinomial Theorem

## Theorem

$$(x_1 + x_2 + x_3 + x_4)^n = \sum_{k_1+k_2+k_3+k_4=n} \binom{n}{k_1, k_2, k_3, k_4} x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4}$$

**Problem** What is the coefficient of  $a^6 b^8 c^6 d^6$  in

$$(4a^3 - 5b + 9c^2 + 7d)^{19}$$

**Answer**  $\binom{19}{2, 8, 3, 6} 4^2 (-5)^8 9^3 7^6$

# The Pigeon Hole Principle

**Old Saying** If you have to put  $n + 1$  pigeons into  $n$  holes, then you must put some two pigeons into the same hole.


**More generally** If you have to put  $mn + 1$  pigeons into  $n$  holes, then you must put some  $m + 1$  pigeons into the same hole.

# The Erdős-Szekeres Theorem

**Theorem** Any sequence of  $m n + 1$  distinct real numbers either contains an increasing subsequence of length  $m + 1$  or a decreasing subsequence of length  $n + 1$ .



# WTT and Paul Erdős (1988?)



# The Erdős-Szekeres Theorem

**Theorem** Any sequence of  $m n + 1$  distinct real numbers either contains an increasing subsequence of length  $m + 1$  or a decreasing subsequence of length  $n + 1$ .

**Example** Here  $m = 3$  and  $n = 5$

2, 3, -5, 0,  $\pi$ , 9, -4, -3, 7, 8, 5, 1, -6, 10, -8, -1

**Problem** Find the longest increasing subsequence and the longest decreasing subsequence.

# Proof of the Erdős-Szekeres Theorem

**Proof** Let the sequence be:  $(a_1, a_2, a_3, \dots, a_t)$  with  $t = mn+1$ . For each  $i$ , place the pigeon  $a_i$  in the pigeon hole  $(inc, dec)$  where  $inc$  is the length of the longest increasing subsequence starting with  $a_i$  and  $dec$  is the length of the longest decreasing subsequence starting with  $a_i$ . Then there are  $mn + 1$  pigeons and only  $mn$  holes. Consider two pigeons  $a_i$  and  $a_j$  which are assigned to the same hole where  $i < j$ . If  $a_i < a_j$ , then  $inc$  value for  $a_i$  is larger than the  $inc$  value for  $a_j$ . If  $a_i > a_j$ , then  $dec$  value for  $a_i$  is larger than the  $dec$  value for  $a_j$ .

# Easy Application

---

**Exercise** Show that if  $S$  is a subset of size 6 from  $\{1, 2, 3, \dots, 9\}$ , then there are two distinct elements of  $S$  whose sum is 10.

# Complexity and Problem Size

**Observation** We typically say that a problem size is  $n$  when the data for a problem can be interpreted as  $n$  packets of information with each packet readable in some constant amount of time. For example, when integers are at most `MAX_INTEGER` in some programming language, we view all integers as being readable in constant time. With this notion in mind, we begin to discuss - quite informally - the running time for algorithms to solve a problem of size  $n$ .

# Running Time

**Concept** An algorithm accepts input data of size  $n$  and then carries out a procedure which takes a total of  $f(n)$  steps. The function  $f(n)$  is called the running time of the algorithm. Typically, it is not possible to determine  $f(n)$  precisely, so it is very important to be able to estimate  $f(n)$  in a reasonably accurate manner.

**Concept** It is also very important to be able to compare two algorithms, one with running time  $f(n)$  and the other with running time  $g(n)$ .

# A Small Catalog of Increasing Functions

**Fact** The following functions all increase and tend to infinity:

$\log^* n$      $\log \log \log n$      $\log \log n$      $\log n$      $n^{.001}$

$\sqrt{n}$      $n$      $n \log n$      $n^{3/2}$      $n^2$      $n^3$      $n^{\log \log n}$

$n^{\log n}$      $2^n$      $(\log n)^n$      $(\sqrt{n})^n$      $n^n$      $2^{2^n}$      $2^{2^{2^n}}$

# The Big-Oh Notation

**Formally** When  $f(n)$  and  $g(n)$  are two functions, the notation  $f = O(g)$ , which is sometimes written  $f(n) = O(g(n))$  means there is a constant  $C > 0$  so that  $f(n) \leq C g(n)$  for all  $n$ .

# The Big-Oh Notation

**Example** Consider two algorithms  $A_1$  and  $A_2$  which have running time, respectively, of  $30n^2 + 150n \log n$  and  $n^3 + n$  respectively. Which one is faster?

Clearly, when the problem size is small, the second one wins, but when the problem size becomes reasonably large, the first one is better. Formally, if we set

$$f(n) = 30n^2 + 150n \log n \quad \text{and} \\ g(n) = n^3 + n$$

In this case, we can write  $f = O(g)$ .

# The Little-Oh Notation

**Formally** When  $f(n)$  and  $g(n)$  are two functions, the notation  $f = o(g)$ , which is sometimes written  $f(n) = o(g(n))$  means that the ratio  $f(n)/g(n)$  tends to 0 as  $n$  tends to infinity, i.e.,

$$\lim_{n \rightarrow \infty} \left( \frac{f(n)}{g(n)} \right) = 0$$

# Applying the Little-Oh Notation

**Example** Compare the two functions:

$$f(n) = 30 n^2 + 150 n \log n \quad \text{and}$$

$$g(n) = n^3 + n$$

When  $n$  is of modest size,  $g(n)$  is smaller, but from some point on  $f(n)$  is smaller. In fact the ratio  $f(n)/g(n)$  goes to 0. In this case, we can write

$$f = o(g).$$

# Revisiting Increasing Functions

**Fact** In each case, if  $f(n)$  and  $g(n)$  are two functions occurring consecutively in this list, then  $f = o(g)$ .

$\log^* n$      $\log \log \log n$      $\log \log n$      $\log n$      $n^{.001}$

$\sqrt{n}$      $n$      $n \log n$      $n^{3/2}$      $n^2$      $n^3$      $n^{\log \log n}$

$n^{\log n}$      $2^n$      $(\log n)^n$      $(\sqrt{n})^n$      $n^n$      $2^{2^n}$      $2^{2^{2^n}}$

# Four Motivating Problems

**Observation** We want to develop a framework for discussing the difficulty of a problem. As examples, given a list  $S$  of  $n$  distinct positive integers, consider the following problems:

1. What is the largest integer in  $S$ ?
2. If  $a$  is the first integer in  $S$ , are there distinct integers  $b$  and  $c$  in  $S$  so that  $a = b + c$ ?
3. Are there three distinct integers  $a, b, c$  in  $S$  with  $a = b + c$ ?
4. Can we solve the fair division problem for  $S$ ?

# Elementary Algorithms

**Example** Finding the largest integer in  $S$  can be done in  $n$  steps where  $n = |S|$  is the problem size.

**Example** If  $a$  is the first number in  $S$ , determining whether there are numbers  $b$  and  $c$  in  $S$  so that  $a = b + c$  can be done in  $\binom{n-1}{2}$  steps.

**Example** Determining whether there are numbers  $a$ ,  $b$  and  $c$  in  $S$  so that  $a = b + c$  can be done in  $\binom{n}{3}$  steps.

# The Fair Division Problem

**Example** Determining whether the fair division problem can be solved for  $S$  can be done in  $n2^n$  steps. Each step consists of choosing a subset  $T$  of  $S$ , and adding all the numbers in  $T$ , adding all the numbers in  $S - T$  and then checking whether the answers are the same. In contrast to the other algorithms, each step takes a formidable amount of time, as  $n - 1$  additions and one comparison have to be made.

# Sorting

**Sorting Problem** You are told that there is an unknown linear order on the integers in  $\{1, 2, \dots, n\}$  and your job is to discover this order by asking questions of the form: Is  $i < j$  in  $L$ ? For example, if  $L = (2, 5, 3, 1, 4)$ , you might discover it by asking:

Is $2 < 1$ in $L$ ?	Answer Yes
Is $4 < 3$ in $L$ ?	Answer No
Is $3 < 1$ in $L$ ?	Answer Yes
Is $2 < 5$ in $L$ ?	Answer Yes
Is $3 < 5$ in $L$ ?	Answer No
Is $1 < 4$ in $L$ ?	Answer Yes

# The UGA Sorting Algorithm

**UGA Sorting Algorithm** For each 2-element subset  $\{i, j\}$  of  $\{1, 2, \dots, n\}$ , ask the question:

Is  $i < j$  in  $L$ ?

So a total of  $\binom{n}{2} = \frac{n(n-1)}{2}$  questions are asked.

Now you have all the information and can easily assemble the linear order  $L$  with these answers.

# Lower Bound on Sorting

**Theorem** In worst case, any sorting algorithm can be forced to ask  $\log_2 n!$  questions.

**Explanation** There are  $n!$  different linear orders (permutations) of  $\{1, 2, \dots, n\}$ . Each time a question is asked, in worst case the number of possible answers is reduced by at most  $\frac{1}{2}$ . So if  $t$  questions are asked, we must have  $2^t \geq n!$

This is equivalent to  $t \geq \log_2 n!$

# The Stirling Approximation

**Theorem** In advanced calculus courses, the following asymptotic formula is proved:

$$\lim_{n \rightarrow \infty} \frac{n!}{\sqrt{2\pi n} \left(\frac{n}{e}\right)^n} = 1$$

It follows that  $\log_2 n! \sim n \log_2 n$ .

**Accordingly** A sorting algorithm is said to be optimal if its running time is  $O(n \log n)$ .

# Some Optimal Sorting Algorithms

---

**Fact** The list of optimal sorting algorithms includes: merge sort, heap sort, introsort, Timsort, Cubesort and Block Sort (there are others).

# An Informal Discussion of Merge Sort

**Concept** To determine an unknown linear order on  $\{1, 2, \dots, n\}$ , first divide the problem into two equal size subproblems. That is, first find the restriction to the subset consisting of the first  $n/2$  integers  $\{1, 2, \dots, n/2\}$ . Then find the restriction to the last  $n/2$  integers  $\{n/2 + 1, n/2 + 2, \dots, n\}$ . Then “merge” the two answers to determine the full linear order.

**Fact** Two sorted lists of size  $n/2$  can be merged in running time  $n$ .

**Fact** Merge sort has a running time  $r(n)$  satisfying the recurrence  $r(n) = 2r(n/2) + n$ . An easy calculation now shows that  $r(n) = O(n \log n)$ .